

An Approach to Optimizing the Water Resources Management Problem in a Fuzzy Environment

El- Saed Ebrahim Ammar & Hamiden Abd Elwahed Khalifa*

El- Saed Ebrahim Ammar, Department of Mathematics, Faculty of Science, Tanta University, Tanta, Egypt

Hamiden Abd Elwahed Khalifa, Department of Operations Research, Institute of Statistical Studies and Research, Cairo University, Giza, Egypt

KEYWORDS

Water resources management;
Fuzzy numbers;
Membership function;
Fuzzy arithmetic;
Fuzzy constraints;
Two stage;
Policy analysis;
Uncertainty.

ABSTRACT

Fully fuzzy linear programming is applied to water resources management due to its close connection with human life, which is considered to be of great importance. This paper investigates the decision-making concerning water resources management under uncertainty based on two-stage stochastic fuzzy linear programming. A solution method for solving the problem with fuzziness in relations is suggested to prove its applicability. The purpose of the method is to generate a set of solutions for water resources planning that helps the decision-maker make a tradeoff between economic efficiency and risk violation of the constraints. Finally, a numerical example is given and is approached by the proposed method.

© 2019 IUST Publication, IJIEPR. Vol. 30, No. 1, All Rights Reserved

1. Introduction

Water Resources Management is an integrating concept for a number of water sub-sectors such as hydropower, water supply and sanitation, irrigation and drainage, and environment (Gasinov and Yenilmez[7]).

The water resources management includes (Jairaj and Vedula[10]):

- The quantitative and qualitative exploration of water resources;
- Water requiring inventory records;
- Measurement and matching of the water resources and water needs (demands) in a special system;
- Decision support depending on the results.

Up to now, fuzzy set theory has been applied to broad fields. Fuzzy set theory introduced by Zadeh [28] creates a model that is set up using

approximately known data. Fuzzy numerical data can be represented by means of fuzzy subsets of the real line, known as fuzzy numbers. For the fuzzy set theory development, we may refer to the papers of Kaufmann [12] and Dubois and Prade [3]. They extended the application of algebraic operations of real numbers to fuzzy numbers by using a fuzzy principle. Fuzzy linear constraints with fuzzy numbers were studied by Dubois and Prade [3]. Lu et al. [16] introduced the definition of an inexact rough interval fuzzy linear programming method and investigated the allocation of generated water to agricultural irrigation system. In the real-world problems, uncertainties may be estimated as intervals. Shaocheng [20] studied two kinds of linear programming with fuzzy numbers called interval numbers and fuzzy number linear programming. Tanaka et al. [22] formulated and proposed a method for solving linear programming with fuzzy coefficients. Wang and Huang [25] developed interactive two-stage stochastic fuzzy programming for managing water resources. They proposed an interactive resolution method

* Corresponding author: *Hamiden Abd Elwahed Khalifa*

Email: hamiden_2008@yahoo.com

Received 28 November 2018; revised 20 February 2019; accepted 05 March 2019

within inexact two-stage stochastic programming. A two-stage optimization framework for planning reservoir operations was proposed by Wang and Adams [23], where hydrologic uncertainty and seasonal reservoir inflows have been modeled as in a periodic Markov process. Through a two-stage dynamic programming approach, a long-term hydrothermal scheduling of multi-reservoir systems was examined by Ferrero et al. [5]. Bellman and Zadeh [1] introduced the concept of a maximizing decision-making problem. Zhao et al. [29] introduced a complete solution set for fuzzy linear programming problems using linear and nonlinear membership functions. For water resources management, Huang and Loucks [9] proposed inexact two-stage stochastic programming. An interactive fuzzy resolution method for solving linear programming problems with fuzzy parameters was proposed by Jimenez et al. [11]. For developing water resources management, a number of optimization techniques were developed (Slowinski [21], Wu et al. [26], Jairaj and Vedula [10], and Maqsood et al. [17]). A model for obtaining an optimal multi-period operation within a multi-reservoir system was developed by Eiger and Shamir [4]. Xu et al. [27] investigated and applied an inexact two-stage fuzzy gradient chance-constrained programming method to the water resources management in Heshui River Basin, Jiangxi Province. To quantify the economic trade-offs when reducing groundwater abstraction to a sustainable level, Martinsen et al. [18] applied a multi-objective multi-temporal deterministic hydro economic optimization approach for this purpose. Fu et al. [6] proposed a two-level symmetric Nash-Harany leader-follower game model to resolve the conflict that arises when different water users compete for a limited water supply. Khalifa [14] studied the water allocation problem using the two-stage fuzzy random programming. An interval-valued fuzzy linear programming method for modeling parameters with high vagueness was represented by Wang et al. [24], Goralczany [8], and Cai et al. [2]. Khalifa and Al-Shabi [15] developed an approach for optimizing the water resources management problem based on the weighting method. This paper aims to introduce and solve the problem of water resources management as two-stage stochastic fuzzy linear programming. The problem is considered by incorporating fuzzy numbers. A solution method for solving the problem with fuzziness in relations is suggested to demonstrate its applicability.

The remainder of the paper is organized as follows. Some preliminaries are given in Section 2. In Section 3, a water resources management problem introduced by Huang and Loucks[9] and Wang and Huang [25] is introduced; hence, the problem is investigated in a fuzzy environment. Characterization of α – fuzzy optimal solution of the problem is presented in Section 4. A solution method for solving the problem is proposed in Section 5. In Section 6, a numerical example is given for illustration. Finally, some concluding remarks are reported in Section 7.

2. Preliminaries

Some basic concepts and results related to fuzzy numbers as well as some of their arithmetic operations, triangular fuzzy numbers, and some of algebraic operations are recalled in this section (Kaufmann and Gupta [13], Sakawa [19] and Zimmermann[30]).

Definition 1.

A fuzzy number \tilde{p} is a mapping:

$\mu_{\tilde{p}} : R \rightarrow [0, 1]$, with the following properties:

- (i) $\mu_{\tilde{p}}(x)$ is an upper semi-continuous membership function;
- (ii) \tilde{p} is a convex set, i. e. , $\mu_{\tilde{p}}(\lambda x + (1 - \lambda)y) \geq \min\{\mu_{\tilde{p}}(x), \mu_{\tilde{p}}(y)\}$ for all $x, y \in R, 0 \leq \lambda \leq 1$;
- (iii) \tilde{p} is normal, i.e., $\exists x_0 \in R$ for which $\mu_{\tilde{p}}(x) = 1$;

Suppose that $(\tilde{p}) = \{x : \mu_{\tilde{p}}(x) > 0\}$ is the support of a fuzzy set \tilde{p} .

Let $F_0(R)$ denote the set of all compact fuzzy numbers on R , that is, for any $g \in F(R)$, g satisfies the following:

1. $\exists x \in R : g(x) = 1$;
2. For any $0 < \alpha \leq 1$, $g_\alpha = [g_\alpha^L, g_\alpha^U]$ is a close interval number on R .

Definition 2.

The α – level set of $\tilde{p} \in F(R), 0 \leq \alpha \leq 1$ is denoted by $(\tilde{p})_\alpha$ and is defined as follows:

$$(\tilde{p})_\alpha = \begin{cases} \{x \in R : \mu_{\tilde{p}}(x) \geq \alpha, 0 < \alpha \leq 1 \\ \text{closure}(\text{support}(\tilde{p})), & \alpha = 0 \end{cases}$$

Definition 3.

A fuzzy number \tilde{A} on R is called triangular fuzzy number, if there exist real a, b, c , and $b, c \geq 0$ such that:

$$\tilde{A}(x) = \begin{cases} \frac{x}{b} + \frac{b-a}{b}, & b-a \leq x \leq a, \\ \frac{-x}{c} + \frac{a+c}{c}, & a \leq x \leq a+c, \\ 0, & \text{elseswhere} \end{cases}$$

Let the triangular fuzzy number denoted by $\tilde{A} = (a, b, c)$ and $F(R)$ be the set of all $L - R$ fuzzy numbers on R .

Definition 4.

$\tilde{A} = (a, b, c)$ is called non-negative triangular fuzzy number if $a \geq 0$.

Definition 5.

Let $\tilde{A} = (a, b, c) \geq \tilde{0}$, $\tilde{B} = (d, e, f) \geq \tilde{0}$, and $x \in R$; the formulas for the addition, subtraction, scalar multiplication, and multiplication can be defined as follows:

1. Addition:

$$\begin{aligned} \tilde{A} \oplus \tilde{B} &= (a, b, c) \oplus (d, e, f) \\ &= (a + d, b + e, c + f). \end{aligned}$$

2. Subtraction:

$$\begin{aligned} \tilde{A} (-) \tilde{B} &= (a, b, c) \oplus (d, e, f) \\ &= (a - f, b - e, c - d). \end{aligned}$$

3. Multiplication:

$$\tilde{A} \otimes \tilde{B} = \begin{cases} (ad, be, cf), & a \geq 0 \\ (af, be, cd), & a < 0, c \geq 0 \\ (af, be, cd), & c < 0 \end{cases}$$

4. Scalar multiplication

$$x \tilde{A} = \begin{cases} (xa, xb, xc), & x \geq 0, \\ (xc, xb, xa), & x < 0. \end{cases}$$

Remark1. Let $\tilde{0} = (0, 0, 0)$ represent a zero triangular fuzzy number.

Remark2. $\tilde{A} \geq \tilde{0}$ if and only if $a \geq 0, a - b \geq 0, a + c \geq 0$.

3. Water Resources Management Problem

In this section, some of the notations needed in the problem formulation are introduced.

3-1. Notations

The following notations are needed in the formulation

f : A benefit of system (\$);

B_j : Net benefit to user j per m^3 of water allocated (\$/ m^3)

(First-stage revenue parameters)

T_j : Allocation target for water that is promised to the user j (m^3)

(First-Stage decision variables)

$E [.]$: Expected value of a random variable;

C_j : Loss to user j per m^3 of water not delivered,

$C_j > NB_j$ (\$/ m^3)

(Second-Stage cost parameters)

S_{jQ} : Shortage of water to user j when the seasonal flow is Q (m^3)

(Second-Stage decision variables))

Q : Total amount of seasonal flow (m^3) (random variables);

δ : Rate of water loss during transportation;

$T_{j\max}$: Maximum allowable allocation amount for user j (m^3);

m : Total number of water users;

i : Water user, $i = 1, 2, 3$, where $i = 1$ for municipality, $i = 2$ for the industrial user, and $i = 3$ for the agricultural sector.

The typical two-stage stochastic programming for the water resources management problem, introduced by Huang and Loucks [9], Wang and Huang [25], is considered.

$$\max f = \sum_{j=1}^n \tilde{B}_j T_j - E \left[\sum_{j=1}^n C_j S_{jQ} \right] \tag{1}$$

subject to

$$\sum_{j=1}^n (T_j - S_{jQ})(1 + \delta) \leq Q, \tag{2}$$

(Water availability constraints)

$$S_{jQ} \leq T_j \leq T_{j\max}; \forall j, \tag{3}$$

(Water-allocation target constraints)

$$S_{jQ} \geq 0; \forall j \tag{4}$$

(Non-negative and technical constraints)
Problems (1)- (4) can be reformulated as in the following form (Huang and Loucks [9]):

$$\max f = \sum_{j=1}^n B_j T_j - \sum_{i=1}^m \sum_{j=1}^n p_i C_j S_{ij} \quad (5)$$

Subject to

$$\sum_{j=1}^n (T_j - S_{ij})(1 + \delta) \leq q_i; \forall i, \quad (6)$$

(Water availability constraints)

$$S_{ij} \leq T_j \leq T_{j_{\max}}; \forall i, j \quad (7)$$

(Water-allocation target constraints)

$$S_{ij} \geq 0; \forall i, j \quad (8)$$

(Non-negative and technical constraints)

where S_{ij} is the amount by which water-allocation target T_j is not met when the seasonal flow is q_i with probability p_i .

$$\max f(S_{ij}, \tilde{B}, \tilde{C}, \tilde{T}) = \sum_{j=1}^n (c_j, b_j, t_j) \otimes (u_j, v_j, w_j) - \sum_{i=1}^m \sum_{j=1}^n p_i ((d_j, e_j, g_j) S_{ij})$$

Subject to

$$S_{ij} \in \tilde{X} = \begin{cases} \sum_{j=1}^n ((u_j, v_j, w_j) - S_{ij})(1 + (\delta_1, \delta_2, \delta_3)) \leq (q_i^1, q_i^2, q_i^3), \forall \\ S_{ij} \leq (u_j, v_j, w_j) (\leq) (T_{j_{\max}}^1, T_{j_{\max}}^2, T_{j_{\max}}^3); \forall, j \\ S_{ij} \geq 0; \forall i, j \end{cases} \quad (10)$$

Based on the arithmetic operations of fuzzy numbers, Problem (10) can be rewritten as follows:

$$\max f(S_{ij}, \tilde{B}, \tilde{C}, \tilde{T}) = \sum_{j=1}^n (c_j, b_j, t_j) \otimes (u_j, v_j, w_j) - \sum_{i=1}^m \sum_{j=1}^n p_i ((d_j, e_j, g_j) \otimes S_{ij})$$

Subject to

3-2. Fuzzy model for water resources management problem

Consider the fuzzy model for problems (5)- (8) as follows:

$$\max f(S_{ij}, \tilde{B}, \tilde{C}, \tilde{T}) = \sum_{j=1}^n \tilde{B}_j \otimes \tilde{T}_j - \sum_{i=1}^m \sum_{j=1}^n p_i (\tilde{C}_j \otimes S_{ij})$$

Subject to

$$S_{ij} \in \tilde{X} = \begin{cases} \sum_{j=1}^n (\tilde{T}_j - S_{ij})(1 + \tilde{\delta}) \leq \tilde{q}_i; \forall \\ S_{ij} (\leq) \tilde{T}_j (\leq) \tilde{T}_{j_{\max}}; \forall, j \\ S_{ij} \geq \tilde{0}; \forall i, j \end{cases} \quad (9)$$

where \tilde{B}_j , \tilde{C}_j , $\tilde{\delta}$, \tilde{q}_i , $\tilde{T}_{j_{\max}}$, and \tilde{T}_j are triangular fuzzy numbers.

Definition 6.

(Optimal fuzzy solution). S_{ij}^* that satisfies the conditions in (9) is called a fuzzy optimization solution.

By using the representation of the fuzzy number as mentioned before, Problem (9) becomes

$$S_{ij} \in X' = \left\{ \begin{array}{l} \sum_{j=1}^n \left((T_j)_\alpha - S_{ij} \right) (1 + \delta_1) \leq q_i^1, i = 1, 2, \dots, m \\ \sum_{j=1}^n \left((T_j)_\alpha - S_{ij} \right) (1 + \delta_1 - \delta_2) \leq (q_i^1 - q_i^2), i = 1, 2, \dots, m \\ \sum_{j=1}^n \left((T_j)_\alpha - S_{ij} \right) (1 + \delta_1 + \delta_3) \leq (q_i^1 + q_i^3), i = 1, 2, \dots, m \\ S_{ij} \leq (T_j)_\alpha \leq T_{j \max}^1, j = 1, 2, \dots, n \\ S_{ij} \leq (T_j)_\alpha \leq T_{j \max}^1 - T_{j \max}^2, j = 1, 2, \dots, n \\ S_{ij} \leq (T_j)_\alpha \leq T_{j \max}^1 + T_{j \max}^3, j = 1, 2, \dots, n \\ S_{ij} \geq 0; \forall i, j \end{array} \right. \quad (11)$$

In problem (11), \tilde{B}_j , \tilde{C}_j , and \tilde{T}_j are the fuzzy variables of R characteristics based on membership functions $\mu_{\tilde{B}_j}$, $\mu_{\tilde{C}_j}$, and $\mu_{\tilde{T}_j}$, respectively.

S^* is an α -fuzzy optimal solution for problem (11) if $\mu(f(S^*, B, C, T) \geq f(S, B, C, T)) \geq \alpha$, for all S_{ij} . (12)

Definition 7.

On the account of the extension principle,

$$\mu(f(S^*, \tilde{B}, \tilde{C}, \tilde{T}) \leq f(S^*, \tilde{B}, \tilde{C}, \tilde{T})) = \sup_{B, C, T} \min(\mu_{\tilde{B}}(B), \mu_{\tilde{C}}(C), \mu_{\tilde{T}}(T)), \quad (13)$$

4. Characterization of α -Fuzzy Optimal Solution Problem (11)

For deducing the α -fuzzy optimal solution for problem (11), let us consider the following α -parametric problem:

Model 1:

$$\max f(S_{ij}, B, C, T) = \sum_{j=1}^n B_j T_j - \sum_{i=1}^m \sum_{j=1}^n p_i (C_j S_{ij})$$

Subject to

$$S_{ij} \in X', B_j \in (\tilde{B}_j)_\alpha, C_j \in (\tilde{C}_j)_\alpha, T_j \in (\tilde{T}_j)_\alpha$$

where $(\tilde{B}_j)_\alpha$, $(\tilde{C}_j)_\alpha$, and $(\tilde{T}_j)_\alpha$ denote the α -cut sets of \tilde{B}_j , \tilde{C}_j , and \tilde{T}_j , respectively. Based on the convexity of the problem, $\mu_{\tilde{B}_j}(B_j)$, $(\tilde{B}_j)_\alpha$; $\mu_{\tilde{C}_j}(C_j)$, $(\tilde{C}_j)_\alpha$; $\mu_{\tilde{T}_j}(T_j)$, $(\tilde{T}_j)_\alpha$ are real intervals that are denoted by $[(\tilde{B}_j(\alpha))^- , (\tilde{B}_j(\alpha))^+]$, $[(\tilde{C}_j(\alpha))^- , (\tilde{C}_j(\alpha))^+]$, and

$[(\tilde{T}_j(\alpha))^- , (\tilde{T}_j(\alpha))^+]$, respectively. Let Ω_α , Δ_α , and Ψ_α be the set of $1 \times n$ matrices $B = (B_j)$, $C = (C_j)$, and $T = (T_j)$ with $B_j \in [(\tilde{B}_j(\alpha))^- , (\tilde{B}_j(\alpha))^+]$, $C_j \in [(\tilde{C}_j(\alpha))^- , (\tilde{C}_j(\alpha))^+]$, and $T_j \in [(\tilde{T}_j(\alpha))^- , (\tilde{T}_j(\alpha))^+]$. It is obvious that problem (14) may be rewritten as follows:

Model. 2.

$$\max f(S_{ij}, B, C, T) = \sum_{j=1}^n B_j T_j - \sum_{i=1}^m \sum_{j=1}^n p_i (C_j S_{ij})$$

Subject to

$$S_{ij} \in X', B_j \in \Omega_\alpha, C_j \in \Delta_\alpha, T_j \in \Psi_\alpha.$$

Definition 8. $S_{ij}^* \in X'$ is called an α -

parametric optimal solution for Model 2, if

$$f(S_{ij}^*, B, C, T) \geq f(S_{ij}, B, C, T); \forall S_{ij} \in X', B_j \in \Omega_\alpha, C_j \in \Delta_\alpha, T_j \in \Psi_\alpha$$

Theorem1. $S_{ij}^* \in \tilde{M}$ is an α – fuzzy optimal solution to problem (9) if and only if $S_{ij}^* \in X'$ is an α – parametric optimal solution to problem Model 2.

Proof. Necessity: Suppose that $(X^*, Y^*) \in \tilde{M}$ is the α – fuzzy optimal solution for problem (2) , and $(X^*, Y^*) \in M'$ is not an α – parametric optimal solution for problem Model 2. Then, there are $S_{ij}^\circ \in \tilde{X}$ and $g \in \Omega_\alpha, l \in \Delta_\alpha, h \in \Psi_\alpha$ such that

$$f(S_{ij}^\circ, g, l, h) \geq f(S_{ij}^*, g, l, h).$$

Since $g \in \Omega_\alpha, l \in \Delta_\alpha, h \in \Psi_\alpha$, we have

$$\mu(f(S_{ij}^\circ, \tilde{B}, \tilde{C}, \tilde{T}) \geq f(S_{ij}^*, \tilde{B}, \tilde{C}, \tilde{T})) \geq \alpha.$$

This contradicts the α – fuzzy optimal solution of $S_{ij}^* \in \tilde{X}$ for problem(9).

Sufficiency: $S_{ij}^* \in X'$ is the α – parametric optimal solution for problem Model 2, and $S_{ij}^* \in \tilde{X}$ is not an α – fuzzy optimal solution for problem(9). Then, there are $S_{ij}^* \in X'$ such that: $\mu(f(S_{ij}^*, \tilde{C}, \tilde{D}) \geq f(S_{ij}, \tilde{C}, \tilde{D})) \geq \alpha$, i.e.,

$$\sup_{B', C, T'} \min(\mu_{\tilde{B}}(B), \mu_{\tilde{C}}(C), \mu_{\tilde{T}}(T)) \geq \alpha, \quad (14)$$

$$B' = \{B \in R^{(1 \times n)} : f(S_{ij}^*, B, C, T) \geq f(S_{ij}, B, C, T)\}, \quad \text{Model. 1-2-2.}$$

$$C' = \{C \in R^{(1 \times n)} : f(S_{ij}^*, B, C, T) \geq f(S_{ij}, B, C, T)\}, \quad \max f^-(S_{ij}, B, C, T) = \sum_{j=1}^n (B_j T_j)^- - \sum_{i=1}^m \sum_{j=1}^n p_i ((C_j)^+ S_{ij})$$

$$T' = \{T \in R^{(1 \times n)} : f(S_{ij}^*, B, C, T) \geq f(S_{ij}, B, C, T)\}$$

For the supremum to exist, there are $\phi \in B', \psi \in C'$, and $\varphi \in T'$ with

$$\min(\mu_{\tilde{B}}(\phi); \mu_{\tilde{C}}(\psi), \mu_{\tilde{T}}(\varphi)) < \alpha, \text{ then}$$

$$\sup_{B', C, T'} \min(\mu_{\tilde{B}}(\phi), \mu_{\tilde{C}}(\psi), \mu_{\tilde{T}}(\varphi)) < \alpha.$$

This contradicts (14). Then, there are $\phi \in B'$, $\psi \in C'$, and $\varphi \in T'$ that satisfy

$$\min(\mu_{\tilde{B}}(\phi); \mu_{\tilde{C}}(\psi), \mu_{\tilde{T}}(\varphi)) \geq \alpha, \text{ i.e.,} \quad (15)$$

$$\phi \in \Omega_\alpha, \psi \in \Delta_\alpha \text{ and } \varphi \notin \Psi_\alpha \quad (16)$$

Based on (14) and (16), the contradiction of the optimality of $S_{ij}^* \in M'$ for problem Model 2 is clear.

Corresponding to Model 2, the following two-level mathematical programming models are structured as follows:

Model. 1-2-1.

$$\max f^+(S_{ij}, B, C, T) = \sum_{j=1}^n (B_j T_j)^+ - \sum_{i=1}^m \sum_{j=1}^n p_i ((C_j)^- S_{ij})$$

Subject to

$$S_{ij} \in X_1 = \begin{cases} \sum_{j=1}^n ((T_j)^- - S_{ij})(1 + \delta_1) \leq q_i^1; \forall \\ \sum_{j=1}^n ((T_j)^- - S_{ij})(1 + \delta_1 - \delta_2) \leq (q_i^1 - q_i^2); \forall \\ \sum_{j=1}^n ((T_j)^- - S_{ij})(1 + \delta_1 + \delta_3) \leq (q_i^1 + q_i^3); \forall \\ S_{ij} \leq (T_j)^- \leq (T_{j\max})^-; \forall, j \\ S_{ij} \leq (T_j)^- \leq (T_{j\max})^-; \forall, j \\ S_{ij} \leq (T_j)^- \leq (T_{j\max})^-; \forall, j \\ B_j \in \Omega_\alpha, C_j \in \Delta_\alpha, T_j \in \Psi_\alpha; \\ S_{ij} \geq 0; \forall i, j \end{cases}$$

Subject to

$$S_{ij} \in X_2 = \begin{cases} \sum_{j=1}^n ((T_j)_\alpha^+ - S_{ij})(1 + \delta_1) \leq q_i^1; \forall \\ \sum_{j=1}^n ((T_j)_\alpha^+ - S_{ij})(1 + \delta_1 - \delta_2) \leq (q_i^1 - q_i^2); \forall \\ \sum_{j=1}^n ((T_j)_\alpha^+ - S_{ij})(1 + \delta_1 + \delta_3) \leq (q_i^1 + q_i^3); \forall \\ S_{ij} \leq (T_j)_\alpha^+ \leq (T_{j\max})_\alpha^+; \forall, j \\ S_{ij} \leq (T_j)_\alpha^+ \leq (T_{j\max})_\alpha^+; \forall, j \\ S_{ij} \leq (T_j)_\alpha^+ \leq (T_{j\max})_\alpha^+; \forall, j \\ B_j \in \Omega_\alpha, C_j \in \Delta_\alpha, T_j \in \Psi_\alpha; \\ S_{ij} \geq 0; \forall i, j \end{cases}$$

Suppose that the α -optimal solutions and the corresponding α -optimum values of Models 2.1 and Model 2.2 are:

$$S_{ij}^{+*}, (f)_\alpha^{+*}; S_{ij}^{-*}, (f)_\alpha^{-*}.$$

5. Solution Method

In this section, a solution procedure for solving the problem (9) is introduced as in the following steps:

- Step1:** Formulate the problem (9),
- Step2:** Transform the problem (9) into the problem (10) and the corresponding problem (11),
- Step3:** Ask the decision-maker to specify α ($0 < \alpha < 1$),
- Step4:** Convert the problem (11) into Model 2,
- Step5:** Use the arithmetic operations of fuzzy numbers to obtain the following two auxiliary models: Model2.1, and Model2.2,

Model. 2-1.

$$\max f^+(S_{ij}, B, C, T) = 644.25 - \left(\begin{array}{l} 70.5S_{11} + 17.25S_{12} + 14.25S_{13} \\ + 117.5S_{21} + 28.75S_{22} + 23.75S_{23} \\ + 47S_{31} + 11.5S_{32} + 9.5S_{33} \end{array} \right)$$

Subject to

Step6: Use any software to obtain the solutions of Model 2.1 and Mode 2.2, hence the optimal fuzzy solution of the problem (9).

6. Numerical Example

Consider the problem introduced by Wang and Huang [25] with triangular fuzzy numbers as: Table1. Economic data (\$/ m³) and seasonal flows (in 10⁶ m³) at different probability levels

Activity		
User		
Municipal ($i = 1$)	Industrial ($i = 2$)	ricultural ($i = 3$)
Maximum allowable	allowable	allocation (\tilde{T}_{\max})
(7, 8, 9)	(7, 8, 9)	(7, 8, 9)
Target of water allocation (\tilde{T}_j)		(1, 2, 3)
(2, 3, 5)	(2, 4, 5)	
Net benefit when water demand is satisfied (\tilde{B}_j)	(95, 100, 110)	(45, 50, 70) (28, 30, 33)
Reduction of the net benefit when demand is not delivered (\tilde{C}_j)	(220, 250, 285)	(55, 60, 90) (45, 50, 75)
Flow level	Probability	Seasonal flow(%)
Low ($i = 1$)	0.3	(2, 3, 4)
Medium ($i = 2$)	0.5	(7, 9, 13)
High ($i = 3$)	0.2	(14, 16, 20)
Water loss ($\tilde{\delta}$)		(0.15, 0.20, 0.40)

$$\begin{aligned}
 S_{11} + S_{12} + S_{13} &\geq 5.2609, \\
 S_{21} + S_{22} + S_{23} &\geq 0.9130, \\
 -S_{31} - S_{32} - S_{33} &\leq 5.1739, \\
 S_{11} + S_{12} + S_{13} &\geq 8.0526, \\
 S_{21} + S_{22} + S_{23} &\geq 9.10526, \\
 S_{31} + S_{32} + S_{33} &\geq 9.10526, \\
 S_{11} + S_{12} + S_{13} &\geq 3.1290, \\
 -S_{21} - S_{22} - S_{23} &\leq 5.9032, \\
 -S_{31} - S_{32} - S_{33} &\leq 14.9355, \\
 S_{ij} &\geq 0; \forall i, j
 \end{aligned}$$

Model. 2-2.

$$\max f^-(S_{ij}, B, C, T) = 352 - \begin{pmatrix} 80.25S_{11} + 22.5S_{12} + 18.75S_{13} \\ +133.75.S_{21} + 37.5S_{22} + 31.25S_{23} \\ +53.5S_{31} + 15S_{32} + 12.5S_{33} \end{pmatrix}$$

Subject to

$$\begin{aligned}
 S_{11} + S_{12} + S_{13} &\geq 5.2609, \\
 S_{21} + S_{22} + S_{23} &\geq 0.9130, \\
 -S_{31} - S_{32} - S_{33} &\leq 5.1739, \\
 S_{11} + S_{12} + S_{13} &\geq 8.0526, \\
 S_{21} + S_{22} + S_{23} &\geq 9.10526, \\
 S_{31} + S_{32} + S_{33} &\geq 9.10526, \\
 S_{11} + S_{12} + S_{13} &\geq 3.1290, \\
 -S_{21} - S_{22} - S_{23} &\leq 5.9032, \\
 -S_{31} - S_{32} - S_{33} &\leq 14.9355, \\
 S_{ij} &\geq 0; \forall i, j
 \end{aligned}$$

Tab. 3. The solution of Model1.2

Optimal policy	Optimum value
$S_{11} = S_{12} = S_{21} = S_{22} = S_{31} = S_{33}$	
$S_{13} = 8.0526$	
$S_{23} = 9.1053$	$f^- = -197.341$
$S_{33} = 9.1053$	

Tab. 2. The solution of Model 1.1

Optimal policy	Optimum value
$S_{11} = S_{12} = S_{21} = S_{23} = S_{31} = S_{33}$	$f^+ = 181.224$
$S_{13} = 8.0526$	
$S_{22} = 9.1053$	
$S_{33} = 9.1053$	

Tab. 4. The fuzzy solution of the problem

Optimal policy	Optimum value
$S_{11} = S_{12} = S_{21} = S_{23} = S_{31} = S_{33}$	$\tilde{f} =$
$S_{13} = 8.0526$	$(0, 0, 308.2354)$
$S_{22} = 9.1053$	
$S_{23} = 9.1053$	
$S_{33} = 9.1053$	

7. Concluding Remarks

In this paper, the water resources management problem was studied under fuzzy environment. Two auxiliary models were obtained from the proposed approach. Each model was solved using Lingo package computer. The advantage of the approach was significant for its use in interactive methods for making any comment by related managers and achieving the solutions logically. Finally, fully fuzzy linear programming for water resources management is recommended while considering minimal S_{ij} by which the water-allocation target, T_j , is not met when the seasonal flow with the probability of p_i is q_i .

Acknowledgment

The author gratefully thanks the anonymous referees for their insightful and constructive comments and suggestions that have led to an improved version of this paper.

References

- [1] Bellman, R. E., and Zadeh, L. A., "Decision making in a fuzzy environment," *Management Science*, Vol. 17, (1970), pp. 141-164.
- [2] Cai, Y. P., Huang, G. H., and Lu, H. W., "I- VFRP: an interval- valued fuzzy robust programming approach for municipal waste- management planning under uncertainty," *Engineering Optimization*, Vol. 41, (2009), pp. 399- 418.
- [3] Dubois, D., and Prade, H., "Possibility Theory: An Approach to Computerized Processing of Uncertainty," Plenum, New York, (1980).
- [4] Eiger, G., and Shamir, U., "Operation of reservoir by stochastic programming," *Engineering Optimization*, Vol. 17, (1991), pp. 293- 312.
- [5] Ferrero, R. W., Rivera, J. F., and Shahidehpour, S. M., "A dynamic programming two- stage algorithm for long- term hydrothermal scheduling of multireservoir systems," *IEEE Transactions on Power Systems*, Vol .13, (1998), pp. 1534- 1540.
- [6] Fu, J., Zhong, P., Zhu, F., Ghen, J., Wu, Y., and Xu, B., "Water resources allocation in transboundary river based on asymmetric Nash- harsanyi leader-follower game model, " *Water*, Vol. 10, (2018), pp. 270- 287.
- [7] Gasinov, N.R., and Yenilmez, K., "Solving fuzzy linear programming with linear membership functions, " *Turkish Journal of Mathematics*, Vol. 26 , (2002), pp. 375- 396.
- [8] Gorzalczany, M. B., "A method of inference in approximate reasoning based on interval- valued fuzzy sets, " *Fuzzy Sets and Systems*, Vol. 21, (1987), pp. 1- 17.
- [9] Huang, H. G., and Loucks, D. P., " An inexact two- stage stochastic programming model for water resources management under uncertainty," *Civil Engineering and Environmental Systems*, Vol. 17, (2000), pp. 95- 118.
- [10] Jairaj, P. G., and Vedula, S., "Multi-reservoir system optimization using fuzzy mathematical programming, " *Water Resources Management*, Vol. 14, (2000), pp. 457- 472.
- [11] Jimenez, M., Arenas, M., Bilbao, A., and Rodriguez, M. V., "Linear programming with fuzzy parameters: An interactive method resolution, " *European Journal of Operational Research*, Vol. 177, (2007), pp. 1599- 1609.
- [12] Kaufmann, A., " Introduction to the Theory of Fuzz Subsets," Vol. I. Academic Press, New York. (1975).
- [13] Kaufmann, A., and Gupta, M. M., " Fuzzy Mathematical Models in Engineering and Management Science," Elsevier Science Publishing Company INC., New York, (1988).
- [14] Khalifa, H.A., "On two- stage fuzzy random programming for water resources management, " *African Journal of Mathematics and Computer Science Research*, Vol.8, (2015), pp. 31- 36.
- [15] Khalifa, H.A., and M.M. Al- Shabi, " On application of two- stage stochastic fully fuzzy linear programming for water resources management optimization," *Journal of Advances in Mathematics and Computer Sciences*, Vol. 29, (2018), pp.1- 11.
- [16] Lu, H., Huang, G., and He, L., " An inexact rough- interval fuzzy linear programming method for generating conjunctive water- allocation strategies to agricultural irrigation systems, " *Applied Mathematical Modelling*, Vol. 35, (2011), pp. 4330- 4340.
- [17] Maqsood, I., Huang, G. H., and Yeomans, H. G., " An interval- parameter fuzzy two- stage stochastic program for water resources management under uncertainty," *European Journal of Operational Research*,

- Vol. 167, (2005), pp. 208- 225.
- [18] Mortinsen, G., Liu, S., Mo, X., Bauer-Gottwen, P., "Optimizing water resources allocation in the Haihe River basin underground water sustainability constraints, "Geophysical Research Abstract, Vol. 20, EGU2018- 12870-1, (2018).
- [19] Sakawa, M., " Fuzzy Sets and Interactive Multi-objective Optimization," New York, USA: Plenum Press, (1993).
- [20] Shaocheng, T., "Interval number and fuzzy number linear programming," Fuzzy Sets and Systems, Vol. 66, (1994), pp. 301-306.
- [21] Slowinski, R.A., " multicriteria fuzzy linear programming method for water supply system development planning, " Fuzzy Sets and Systems, Vol. 19, (1986), pp. 217- 237.
- [22] Tanaka, H., Tchihashi, H., and Asai, K., "A formulation of fuzzy linear programming problem based on comparison of fuzzy number, " Control of Cybernetics, Vol. 13, (1984), pp.184- 194.
- [23] Wang, D., and Adams, B.J., "Optimization of real- time reservoir operations with Markov decision processes,". Water Resources Research, Vol. 22, (1986), pp. 345- 352.
- [24] Wang, L.Z., Fang, L., and Hipel, K.W., " Water resources allocations: Cooperative game theoretic approach programming for resources management." Journal of Environmental Informatics, Vol. 2, (2003) pp. 11- 22.
- [25] Wang, S., and Huang, H. G., "Interactive two- stage stochastic fuzzy programming for resources management,". Journal of Environmental Management, Vol. 92, (2011), pp. 1986- 1995.
- [26] Wu, S. M., Huang, G. H., and Guo, H. C., "An interactive inexact- fuzzy approach for multi-objective planning of water resource systems,". Water Science and Technology, Vol. 36, (1997), pp. 235- 242.
- [27] Xu, J., Huang, G., Li, Z., and Chen, J., "A two- stage fuzzy chance- constrained water management model., " Environmental Science and Pollution Research, Vol. 24, (2017), pp. 12437-12454.
- [28] Zadeh, L. A., "Fuzzy sets," Information Control, Vol. 8, (1965), pp. 338- 353.
- [29] Zhao, R., Govind, R., and Fan, G., "The complete decision set of the generalized symmetrical fuzzy linear programming problem, " Fuzzy Sets and Systems, Vol. 51, (1992), pp. 53-65.
- [30] Zimmermann, H.J, "Fuzzy sets Theory and Applications," Kluwer- Nijhoff. Boston, MA, (1990).

Follow This Article at The Following Site:

Khalifa H. A. E., Ammar E. S. E., An Approach to Optimizing the Water Resources Management Problem in a Fuzzy Environment. IJIEPR. 2019; 30 (1) :1-10
URL: <http://ijiepr.iust.ac.ir/article-1-868-en.html>

